

The INChO Examination Board when it met in March 2009 reconsidered the solutions to INChO 2009. The answer key to Q.No 7.8 was revised and the answer scripts of **all** students were reassessed for this question. In light of the above, all students whose marks have changed are being sent fresh performance cards. The OCSC list for chemistry which is displayed on the website has been prepared taking into account this change of marks.

**Equivalent Solutions may exist.****Problem 1**

---

**Hydrogen atom**

1.1

1.2

1.3

1.4

1.5 Ans: 16

1.6






1.7 a) Ans : 4

b)



1.8

Problem 2

Nitrogen containing compounds






2.5 i) **D is more basic than E**




2.7 i)  $C_{16}H_{13}NO_4$


2.8




2.9




2.10



2.11





2.12



## Problem 3

## Chemistry of silicon

3.1


i) CO<sub>2</sub>ii) SiO<sub>2</sub>

3.2. b) carbon has small size and forms a  $\pi$  bond with good overlap whereas silicon has larger size hence has a poor  $\pi$  overlap

3.3 b) silicon has larger atomic size than carbon

c) silicon has 3d orbitals which form an  $sp^3d^2$  hybrid orbitals

3.4




3.5 c) bond strength

3.6




3.7




3.8 b) impurities are more soluble in liquid phase than in solid

3.9 c) silicon has low lying unoccupied orbitals


3.10



3.11



3.12




Problem 4

Natural compounds and intermediates


4.1 b) 256

4.2




4.3


a)




b)




c) i)







ii)








4.4




4.5



4.6



4.7







Problem 5

Chemical kinetics

5.1



5.2



5.3

a) 
$$\frac{d[A]}{dt} = -k$$

b) 
$$[A] = [A]_0 - kt$$

c) 
$$t_{0.5} = \frac{[A]_0}{2k}$$

5.4

a) 0

b) 0.11 torr/s

5.5

a) 1

b)  $2.3 \times 10^{-5} \text{ s}^{-1}$

5.6

3240 years

(i)  $3.05 \times 10^{-7} \text{ mol}$

ii)  $6.3 \times 10^{16}$

iii)  $1.9 \times 10^{17}$

iv)  $6.2 \times 10^{23} \text{ mol}^{-1}$

Problem 6

**A. Kinetic theory of gases and Gas Laws**

6.1.	i)	<b>Curve</b>	<b>Temperature</b>
		Curve a	100K
		Curve b	300K
		Curve c	700 K

ii) True

6.2	<b>Curve</b>	<b>Gas</b>
	Curve a	$H_2$
	Curve b	$CH_4$
	Curve c	$NH_3$
	Curve d	Ar

6.3 iii)  $H_2 < CH_4 < NH_3$

6.4 2 atm


6.5 134.7 J

6.6 6.7%

**B.**  
6.7 Solid/vapor

6.8 T= 217 K and P = 5.2 atm

6.9




6.10 a) All the three phases are in equilibrium

b) Molar Gibbs energy for the three phases is the same

6.11 a) increase

6.12 Single

6.13



6.14 a) K decreases as the temperature rises

## Problem 7

## Acid-Base chemistry


## Part A

7.1 a)  b)  c) 7.2 7.3 

## Part B

7.4 
$$\text{CO}_3^{2-} + \text{H}^+ \rightarrow \text{HCO}_3^-$$
$$\text{HCO}_3^- + \text{H}^+ \rightarrow \text{H}_2\text{CO}_3 \quad \text{or} \quad \text{H}_2\text{O} + \text{CO}_2$$
7.5 a)  $1.104 \times 10^{-3}$  mol HCl  
b)  $3.944 \times 10^{-3}$  mol HCl7.6 46.8 %  $\text{Na}_2\text{CO}_3$   
29.1 %  $\text{NaHCO}_3$

7.7



- 7.8
- The total volume of HCl required to reach the 2<sup>nd</sup> end point is twice that of the first one
  - Number of moles of  $\text{CO}_3^{2-}$  is equal to the number of moles of  $\text{HCO}_3^-$  at some point on this curve
  - Number of moles of  $\text{HCO}_3^-$  is equal to twice the number of moles of  $\text{CO}_3^{2-}$  at some point on this curve